



**Mi Dr hospital.**

C. Salvador Novo 1060, Zona Rio, Tijuana, Baja California.

Telephone: 011-52664-682-8880.

e-mail: dreliasortiz@hotmail.com

**MEDICAL REPORT  
GASTRIC SLEEVE SURGERY**

TO WHOM IT MAY CONCERN:

Patient: DEBRA STEIMEN.

Diagnosis: OBESITY.

Surgery date: FEB 01, 2014.

Surgery Type: LAPAROSCOPIC GASTRIC SLEEVE, REVISION.

1.- PRESURGICAL DIAGNOSIS: PREVIOUSLY LAP BAND SURGERY, LAP BAND REMOVAL AND GASTRIC SLEEVE SURGERY.

2.- POSTSURGICAL DIAGNOSIS: GASTRIC PLICATION VS GASTRIC SLEEVE SURGERY.

3.- SURGERY SCHEDULED: LAPAROSCOPIC GASTRIC SLEEVE SURGERY VS GASTRIC BY PASS.

4.- SURGERY PERFORMED: LAPAROSCOPIC GASTRIC SLEEVE SURGERY.

5.- SURGEON: DR ELIAS ORTIZ.

6.- ANESTHESIOLOGIST: DR MIGUEL A. PEREZ MALAGON.

7.- ANESTHESIA: GENERAL

8.- FINDINGS: ADHESIONS CAUSED FROM PAST SURGERIES, STOMACH DECREASED IN SIZE, THE PRESENCE OF STAPLES WAS NOT OBSERVED, SO DIFFICULT TO DETERMINE THE TYPE OF BARIATRIC SURGERY PERFORMED PREVIOUSLY

9.- TECHNICAL: UNDER GENERAL ANESTHESIA AND WITH THE PATIENT LYING ON THE OPERATION TABLE, THE LAPAROSCOPIC PORTS WERE INTRODUCED UNDER DIRECT VISION.

AFTER REMOVING ALL THE ADHESIONS FROM THE PAST SURGERIES THE STOMACH WAS DIVIDED BY FIRING STAPLES.

DURING THIS PROCESS THE SIZE OF THE GASTRIC REMNANT WAS CALIBRATED USING A 33 Fr BOUGIE. AFTER PLACEMENT OF THE BOUGIE, ADDITIONAL STAPLES WERE FIRED AT AN ANGLE. THIS COMPLETED THE CREATION OF THE SLEEVE GASTRECTOMY, A NONABSORBABLE SUTURE WAS THEN APPLIED TO REINFORCE THE STAPLE LINE.

NEXT THE STOMACH WAS EXTRACTED AND A DRAIN WAS PLACED.

THE SURGERY ENDED WITHOUT COMPLICATIONS..

10.- BLEEDING:

11.- COMPLICATIONS:

12.- INCIDENTAL:

13.- ACCOUNT COMPLETE MATERIAL

THE DAY AFTER THE SURGERY A CHECK-UP WAS CONDUCTED. (FLUOROSCOPY WITH CONTRAST SWALLOW).

THE PROJECTION/FILM WAS PRINTED AND GIVEN TO THE PATIENT.

THE PATIENT WAS DISCHARGED WITHOUT SURGICAL OR INTRA ABDOMINAL WOUND COMPLICATIONS. THE PATIENT TOLERATED CLEAR LIQUIDS AND WAS AMBULATORY.

THE FOLLOWING MEDICATIONS WERE PRESCRIBED:

**CIPROFLOXACIN (ANTIBIOTIC)** THESE TABLETS MUST BE CRUSHED BEFORE TAKING THEM. TAKE EVERY 12 HOURS FOR 6 DAYS.

**SUBLINGUAL KETOROLAC**.FOR PAIN. ONE TABLET TO BE TAKEN EVERY 6 HOURS.

A handwritten signature in blue ink, appearing to read 'E. Ortiz'.

---

Dr. Elias Ortiz - Gomez  
Bariatric Surgeon